

A framework for European soil monitoring

Mark Kibblewhite^A, D Arrouays^B, X Morvan^B, R Baritz^C, E Eberhardt^C, S Huber^D, G Prokop^D, RJA. Jones^A, FJA. Verheijen^D, E Micheli^E, T Szegi^E, L Montaranella^F and M Stephens^A

^A National Soil Resources Institute, Natural Resources Department, Cranfield University, Cranfield, Bedfordshire MK43 0AL, United Kingdom, E-mail: m.kibblewhite@cranfield.ac.uk

^B INRA, Centre de Recherches d'Orléans, US 1106, BP 20619 Ardon, 41566 Olivet, France,

^C Bundesanstalt für Geowissenschaften und Rohstoffe, Stilleweg 2, Hannover, 30655, Germany,

^D Umweltbundesamt GMBH, Spittelauer Laende 5, Wien, 1090, Austria,

^E Szent Istvan Egyetem, pater Karoly u. 1., Godollo, 2100, Hungary

^F Joint Research Centre, Ispra, Italy

Abstract

The ENVASSO project has developed a framework for monitoring European soils. 27 indicators were selected for erosion, organic matter decline, contamination, compaction, salinisation, decline in biodiversity, soil sealing, landslides and desertification. A monitoring network with a density of 1 site per 300 km² covers most soil type and land use combinations. 20 indicators were qualified for implementation, covering soil erosion by water, decline in soil organic matter, soil contamination, soil sealing, compaction, salinisation and desertification. Methods for monitoring wind erosion, tillage erosion and carbon stocks in peat soils were found to be inadequate. A tiered approach to implementation of soil monitoring is recommended.

Key words

Monitoring, Indicators.

Introduction

Serious threats to soil throughout Europe have been identified (European Commission 2002, 2006). However, more evidence is needed to support stronger soil protection policy and to target and monitor its implementation. This requires a European soil monitoring network. Some national networks exist (Jones *et al.* 2005) but many have not been re-sampled. The Environmental Assessment of Soil for Monitoring (ENVASSO) project (Kibblewhite *et al.* 2008) aimed to define a European soil monitoring system and describe its potential implementation.

Methods

Indicator selection (Huber *et al.* 2008) was based on international recommendations (OECD 2003). 188 issues and 290 indicators were evaluated. The principal criteria were relevance, methodological soundness, measurability and policy relevance. Priority indicators were chosen (one for each of three key issues per threat) from an initial selection of 27 key issues and 60 indicators. Where possible, performance criteria (e.g. minimum detectable change, background values and indicator thresholds) were defined (Arrouays *et al.* 2008). Methods for within-plot sampling and for parameter measurement were documented (Jones *et al.* 2008). Geo-referenced information was collated on existing soil monitoring networks and inventories that could be re-sampled (Arrouays *et al.* 2008). The number of additional sites needed to adequately monitor soil type and land use combinations was estimated (Morvan *et al.* 2008). Data management requirements were defined (Baritz *et al.* 2008) and a prototype database system (SoDa) constructed with web-based inter-operability. Procedures for estimating 22 indicators were tested in 28 pilot studies (Micheli *et al.* 2008) covering representative regions and land uses.

Results

Of the 27 priority indicators, 20 were qualified (Table 1) but measurement methods are not available or were considered inadequate for 7 others. Better methods are required for continental scale estimation of wind and tillage erosion. Estimation of peat stocks requires reliable methods for estimating the distribution of peat depths and to account for variability in peat composition. A minimum spatial density of 1 site per 300 km² is recommended (Morvan *et al.* 2008), which approximates to a regular grid with sampling sites set at nodes 16 to

17 km apart. This is representative of most combinations of soil types and land uses. The site area should be (Arrouays *et al.* 2008) between 100 m² and 1 ha with homogeneous soil profile development. A minimum of 4 and preferably between 10 and 100 sub-samples fixed depth sub-samples should be taken, depending on the site area and soil profile variation. Pedogenic horizon sampling adjacent to the sample area usefully supports site characterization. The time interval between sampling events needs to be long enough to allow for changes that can be detected within measurement errors. Analysis (Arrouays *et al.* 2008; Morvan *et al.* 2008; Saby *et al.* 2008) indicates that a minimum interval of 10 years is required.

Table 1. Threats, issues and indicators

Threat / Issue	Indicator
Qualified ENVASSO indicators	
Soil erosion	
Water erosion	Estimated soil loss by rill, inter-rill, and sheet erosion
Decline in soil organic matter	
Soil organic matter status	Topsoil organic carbon content (measured) Soil organic carbon stocks (measured)
Soil contamination	
Diffuse contamination	Heavy metal contents in soils Critical load exceedance by S and N
Local soil contamination	Progress in management of contaminated sites
Soil sealing	
Soil sealing	Sealed area
Land consumption	Land take [to urban and infrastructural development]
Brownfield re-development	New settlement area established on previously developed land
Soil compaction	
Compaction, structural degradation	Density Air-filled pore volume at specified suction
Causes of compaction	Vulnerability to compaction
Decline in soil biodiversity	
Species diversity	Earthworm diversity and biomass Collembola diversity
Soil microbial respiration	Microbial respiration
Soil salinisation	
	Salt profile Exchangeable sodium percentage Potential salt sources
Desertification	
	Land area at risk of desertification Land area burnt by wildfire
Non-qualified ENVASSO indicators	
Soil erosion	
Wind erosion	Estimated soil loss by wind erosion
Tillage erosion	Estimated soil loss by tillage erosion
Decline in soil organic matter	
	Peat stock
Landslides	
	Occurrence of landslide activity Volume/mass of displaced material Landslide hazard assessment
Desertification	
	Soil organic carbon content in desertified land

Discussion

Two types of indicators are implemented more easily. Group A are those for which there are existing networks i.e. *topsoil organic carbon contents, heavy metal contents in soils, critical load exceedance by sulphur and nitrogen, salt profile, exchangeable sodium percentage and potential salt sources*. Group B rely on existing remote-sensed data and / or spatial information i.e. *estimated soil loss by rill, inter-rill and sheet erosion, sealed area, land take, vulnerability to compaction, land area at risk of desertification and land area burnt by wild fire*. These groups provide a basis for monitoring soil erosion, decline in soil organic matter, soil contamination, soil sealing, compaction, salinisation and desertification. Some indicators require inventories that do not exist in all

regions or lack harmonization i.e. *progress in the management of contaminated land, new settlement area established on previously developed land* and *occurrence of landslide activity*. Other indicators are compromised by uncertainties in relating site measurements to estimates over space and time, these are *soil density, air capacity, earthworm diversity, Collembola diversity* and *soil microbial respiration*.

A two-tiered approach to implementation is recommended. The first tier should establish a network with a site density of 1 per 300 km², for estimation of indicators in groups A and B. The second tier should be a sub-set of first tier sites with more extended and intensive monitoring, addressing requirements that cannot be implemented feasibly at the first tier, or for which fewer sites are needed. Examples are (1) when measurement procedures are too demanding for general implementation (e.g. some biological, gaseous flux and physical measurements, including measurement of soil erosion), (2) where intensive sampling is needed to describe soil processes to interpret indicator levels and trends (e.g. detailed assessment of sub-soil and lower horizons, or connectivity to landscape processes such as catchment inputs and outputs), (3) special investigations of error sources (e.g. intensive collection and testing of sub-samples to determine an optimum number for application in the first tier), or (4) when performing proficiency exercises to assess variability associated with different field teams (e.g. estimates of stone contents and texture). The second tier network could also provide reference sites for soil typological units.

Conclusions

There is a sufficient density of existing sites for continental soil monitoring over much of the European Union and the number of new sites required is relatively limited and confined to a few member states. There is an adequate technical basis to implement a successful monitoring system for a majority of the threats to soil resources. Current methods are inadequate for assessment of carbon stocks in peat soils, wind erosion and tillage erosion. The ENVASSO manual of procedures and protocols (Jones *et al.* 2008) is a valuable reference for future soil monitoring.

Acknowledgements

This work was conducted under the European 6th Framework Programme of Research ENVASSO Project Contract no. 022173. The financial support of the European Commission is gratefully acknowledged. Some 150 scientists participated in the ENVASSO project and the contribution of each and every one of them is acknowledged with gratitude.

References

- Arrouays D, Morvan X, Saby NPA, Le Bas C, Bellamy PH, Berenyi Uveges J, Freudenschuß A, Jones AR, Jones RJA, Kibblewhite MG, Simota C, Verdoot A, Verheijen FGA (2008) Environmental Assessment of Soil for Monitoring. Volume IIa Inventory and Monitoring. EUR 23490 EN/2A Office of the Official Publications of the European Communities, Luxembourg, 88 pp. DOI 10.2788/93524
- Baritz R, Eberhardt E, Van Liedekerke MH, Panagos P (2008) Environmental Assessment of Soil for Monitoring: Volume III Database Design and Selection. EUR 23490 EN/3 Office for the Official Publications of the European Communities, Luxembourg, 125 pp. DOI 10.2788/93697.
- European Commission (2002) Communication of 16 April 2002 from the Commission to the Council, the European Parliament, the Economic and Social Committee and the Committee of the Regions: Towards a Thematic Strategy for Soil Protection. COM 2002, 179 final.
- European Commission (2006) Communication of 22 September 2006 from the Commission to the Council, the European Parliament, the Economic and Social Committee and the Committee of the Regions: Thematic Strategy for Soil Protection. COM 2006, 231final.
- Huber S, Prokop G., Arrouays D., Banko G., Bispo A., Jones RJA., Kibblewhite MG., Lexer W, Möller A, Rickson RJ, Shishkov T, Stephens M, Toth G., Van den Akker JJH, Varallyay G, Verheijen FGA, Jones AR (2008) Environmental Assessment of Soil for Monitoring: Volume I Indicators and Criteria. EUR 23490 EN/1 Office for the Official Publications of the European Communities, Luxembourg, 339 pp. DOI 10.2788/93515

Jones RJA, Verheijen FGA, Reuter HI, Jones AR (2008) Environmental Assessment of Soil for Monitoring:

- Volume V Procedures and Protocols. EUR 23490 EN/5 Office for the Official Publications of the European Communities, Luxembourg, 165 pp, DOI 10.27788/94366
- Kibblewhite MG, Jones RGA, Montaranella L, Baritz R, Huber S, Arrouays D, Micheli E, Stephens M (2008) Environmental Assessment of Soil for Monitoring: Volume VI Soil Monitoring System for Europe. EUR 23490 EN/6 Office for the Official Publications of the European Communities, Luxembourg, 72 pp
- Micheli E, Bialousz S, Bispo A, Boixedera J, Jones AR, Kibblewhite MG, Kolev N, Kosmas C, Lilja H, Malucelli F, Rubio JL, Stephens M (2008) Environmental Assessment of Soil for Monitoring: Volume IVa Prototype Evaluation Report. EUR 23490 EN/4A Office for the Official Publications of the European Communities, Luxembourg, 96pp. DOI 10.2788/93772
- Morvan X, Saby NPA, Arrouays D, Le Bas C, Jones RJA, Verheijen FGA, Bellamy PH, Stephens M, Kibblewhite MG (2008) Soil monitoring in Europe: a review of existing systems and requirements of harmonization. *Science of the Total Environment* **391**, 1-12.
- OECD (2003) OECD environmental indicators, development, measurement and use. Reference paper, pp 1-37.
- Saby NPA, Bellamy PH, Morvan X, Arrouays D, Jones RJA, Verheijen FGA, Kibblewhite MG, Verdoot AYi, Úveges JB, Freudenschuß A, Simota C (2008) Will European soil-monitoring networks be able to detect changes in topsoil organic carbon? *Global Change Biology* **14**, 1-11.